

MRI-PHY/P981063
 hep-ph/9810339

Semileptonic charm decay as a test for the spectator model in the B_c -meson

Debajyoti Choudhury*, Anirban Kundu† and Biswarup Mukhopadhyaya‡

*Mehta Research Institute,
 Chhatnag Road, Jhusi, Allahabad - 211 019, India*

Abstract

The b and the c quark compete with each other in decays of the B_c -meson. We emphasize the need of obtaining reliable signals of $c \rightarrow s$ in order to assess the merits of the spectator approximation in calculating the relative strengths of the two types of decays. This, we argue, can be done by considering the decay $B_c \rightarrow B_s(B_s^*)l\bar{\nu}_l$, followed by semileptonic decays of the B_s . We suggest looking for like-sign dileptons together with a D_s at the BTeV or LHC-B experiments, and show that such signals can be made background-free by suitable event selection.

*Electronic address: debchou@mri.ernet.in

†Electronic address: akundu@mri.ernet.in

‡Electronic address: biswarup@mri.ernet.in

The recent discovery of the B_c -meson, consisting of a $\bar{b}$ and a c quark, opens up rather interesting possibilities whose test may confront one with some ill-understood aspects of heavy flavour physics [1]. So far, the Collider Detector at Fermilab (CDF) have confirmed observations of the B_c [2]. In addition, the ALEPH, OPAL and DELPHI collaborations at the Large Electron Positron (LEP) collider at CERN have reported candidates for a similar designation [3–5]. The decay channel utilised in all these searches is one where the $\bar{b}$ -quark decays first into a $\bar{c}$, producing a J/ψ , either semi-leptonically or with one or more pions. This is because the J/ψ is easily identified through the decay into two leptons, and the additional leptonic and pionic tracks passing close to the decay vertex provide a viable reconstruction of the B_c [6]. The best fit for the B_c mass obtained from this channel also shows a fair agreement with that predicted by potential model calculations [7].

However, even a naive estimate of the lifetimes of the b and the c quark shows that, in the B_c meson, the charm decay may take precedence over the bottom decay. This is due to the fact that, unlike $b \rightarrow c$, the decay $c \rightarrow s$ is not suppressed by any off-diagonal element of the Cabibbo-Kobayashi-Maskawa (CKM) matrix. Such a suppression tends to offset the effect of the mass of the b -quark. The annihilation channels of B_c are also important [8]. Both these types of decays have already been studied in different theoretical approaches. To be more specific, the semileptonic channels $B_c \rightarrow J/\psi l \bar{\nu}_l$ and $B_c \rightarrow B_s^{(*)} l \bar{\nu}_l$ have been studied using the spectator approximation and different types of form-factors, *e.g.*, BSW [9], ISGW [10], ACCMM [11] etc. No definite conclusion can yet be drawn, however, regarding the preferability of a non-relativistic potential [12] to a relativistic one, when a c decays into an s , while both of them interact with the b -quark all along [13]. It has also been pointed out that the massive spectator for the latter mode significantly affects the phase space that would have been otherwise available with a light spectator [14]. Summing everything up, one can say that the decay widths for the two semileptonic modes might as well be just comparable.

The above remarks show that there is still considerable scope for improving our understanding of the B_c -system where two heavy quarks compete with each other in the decay processes and at the same time influence the transverse motions of each other. In order to clearly establish the validity of a given theoretical picture, it is important at this stage to experimentally identify signatures of decays induced by $c \rightarrow s$. In this note, we suggest ways of extracting the corresponding final states in the semileptonic decay

$$B_c \longrightarrow B_s^{(*)} l \bar{\nu}_l . \quad (1)$$

One of the reasons for concentrating on semileptonic decays is to ensure that our basic question does not get swamped by additional theoretical uncertainties concerning, for example, the legitimacy of factorisation. Thus, our numbers will depend only on the parametrization of the semileptonic decays using spectator approximation, and any observed deviation from these numbers will mean significant non-spectator physics contribution.

As will be clear from the discussion below, it is profitable for us to consider final states which are again produced through semileptonic decays of the B_s . A realistic testing ground for such decays will be provided by experiments dedicated to B-studies at hadronic colliders, typical examples being the BTeV and LHC-B experiments [15]. Depending on the integrated luminosity achieved ultimately, these machines can be expected to produce $10^{11} - 10^{12}$ $b\bar{b}$ -pairs per year. Thus, with a fragmentation ratio of 3.8×10^{-4} (5.4×10^{-4}) for the B_c (B_c^*) meson¹ [16], approximately $10^8 - 10^9$ B_c s can be available according to a rather conservative estimate. One

¹ B_c^* decays almost entirely to B_c , so these two numbers just add up to give the total fragmentation ratio.

may expect a higher yield if one includes the possibility of having a B_c on one side and B -meson of a different type on the other [17].

The decay $B_c \rightarrow B_s^{(*)} l \bar{\nu}_l$ is followed by the B_s decaying into one of its allowed channels. (The B_s^* decays almost entirely into a B_s .) These include final states consisting of $D_s \pi$, $D_s 3\pi$ as well as the semileptonic channel $D_s^{(*)} l \nu_l$. In this letter, we propose to utilise the semileptonic channel alone. Despite suffering from a small branching fraction, this has a unique advantage. Because of the high rate of $B_s - \bar{B}_s$ oscillation, both set of final states $D_s^{+(*)} l^- \bar{\nu}_l$ and $D_s^{-(*)} l^+ \nu_l$ are equally probable. Thus in approximately half the cases one expects like-sign dileptons together with a D_s (including one that may have a D_s^* as its source). This kind of a signal is relatively background-free compared to both the pionic final states and those comprising unlike-sign dileptons. Sources that can fake such signals are limited to those which have two-level neutral particles at the last step of a cascade. The most prominent example of such backgrounds is the decay chain $B^0 \rightarrow D_s^{(*)} D^- \rightarrow D_s K^0 l^- \nu$, following which the K^0 can give rise to leptons of either sign. We shall show that they can be rather easily removed by kinematic cuts.

Calculation of the rates for the exclusive semileptonic decays is straightforward in a form-factor approach. The hadronic matrix elements relevant for the decay of a generic pseudoscalar meson P_0 may be parametrized as

$$\begin{aligned} \langle P(k) | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | P_0(p) \rangle &= f_+ (p + k)_\mu + f_- q_\mu \\ \langle V(k, \epsilon) | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | P_0(p) \rangle &= \frac{2V_{P_0 V}}{m_0 + m_V} \epsilon_{\mu\nu\alpha\beta} \epsilon^{*\nu} p^\alpha k^\beta \\ &+ i \left[(m_0 + m_V) A_{P_0 V}^{(1)} \epsilon_\mu^* \right. \\ &\quad \left. - \epsilon^* \cdot p \left\{ \frac{A_{P_0 V}^{(2)}}{m_0 + m_V} (k + p)_\mu + 2 \frac{m_V}{q^2} [A_{P_0 V}^{(3)} - A_{P_0 V}^{(0)}] q_\mu \right\} \right] \end{aligned} \quad (2)$$

In eq.(2), $q \equiv p - k$ and P and V correspond to generic pseudoscalar and vector mesons respectively. The value of the form-factors $f_\pm$, V and $A^{(i)}$ and their q^2 -dependence can be calculated within a specific model. For the rest of our analysis, we shall use the results obtained within the Bauer-Stech-Wirbel (BSW) formalism [18] which we compile in Table 1. We have checked that our results are insensitive to, say, a 20% variation of the form-factors, including that arising from the uncertainty in the parameter ω in the BSW framework.

In the remainder of our analysis we shall assume that the pions and/or photons resulting from the decay of an excited state(s) to the corresponding ground state will remain untagged. Since the signal consists of a D_s together with a pair of like-sign dileptons, the relevant backgrounds emanate from processes that include these particles in the final state apart from other undetected ones. The most important processes² in this context are

$$B_d^0 \longrightarrow D_s D^- \longrightarrow D_s K^0 l^- \nu, \quad (3a)$$

and

$$B_d^0 \longrightarrow D_s D^- \longrightarrow D_s K^{*0} l^- \nu \longrightarrow D_s K^0 \pi l^- \nu. \quad (3b)$$

²There also exist additional channels with intermediate D_s^* and/or D^{*-} states cascading into the respective ground states. Since the phase space distributions are quite akin to those above, these channels cannot be distinguished unless the accompanying photon(s) are detectable. We shall assume that this is not the case and add on such contributions to those of eqs.(3a) and (3b).

Mode	m_b	m_c	m_s	ω	h_0	h_V	h_{A_1}	h_{A_2}
$B_c \rightarrow B_s$	4.9	1.7	0.55	0.4	0.403	—	—	—
				0.5	0.588	—	—	—
	5.0	1.5	0.5	0.4	0.432	—	—	—
				0.5	0.617	—	—	—
$B_c \rightarrow B_s^*$	4.9	1.7	0.55	0.4	—	3.367	0.487	1.155
				0.5	—	4.794	0.693	1.697
	5.0	1.5	0.5	0.4	—	3.527	0.521	1.246
				0.5	—	4.905	0.725	1.765

Table 1: *Form-factors relevant to the decays of eq.(1) as calculated within the BSW formalism [18]. The values are shown for $q^2 = 0$ only. The relevant $c\bar{s}$ poles are: 2.60 GeV (f^+), 2.11 GeV (V_{P_0V}), 2.53 GeV ($A_{P_0V}^{(1,2)}$). The other formfactors do not contribute in the limit of vanishing lepton masses. All masses and ω are in GeVs. All other form-factors are obtained from Ref. [18].*

The K^0 has a 50% probability each of being in the K_L or the K_S state. With the K_S having a negligibly small semileptonic branching ratio, it is the K_L which can decay (BR $\simeq 65\%$) into $\pi l \nu_l$ ($l = e, \mu$), l^+ and l^- being produced with equal probability.

As $m(D_s) - m(K^{*0})$ is considerably smaller than $m(D_s) - m(K^0)$, lepton from the D_s decay will have significantly different energy spectra in the two cases. Similarly, the leptons in the signal events will have an entirely different profile. We aim next to quantify this difference in a frame-independent manner.

The first kinematic distribution on which we focus our attention is $\Gamma^{-1} d\Gamma/dm_{\ell\ell}$ where $m_{\ell\ell}$ is the invariant mass of the dilepton system. Note that this is a Lorentz invariant distribution and is unaffected by the extent to which the parent B_c s are boosted. For the signal, this distribution shows a peak in the neighbourhood of 1 GeV (see Fig. 1), and has little dependence on the form-factors. The figure clearly shows that the background peaks at a considerably lower invariant mass compared to the signals. This is because each of the two leptons in the background process is produced at a later stage of the cascade, thereby suffering a degradation that lowers the invariant mass.

It must be borne in mind though, that Fig. 1 exhibits only the normalized distributions. However, at a hadronic machine, the production rate for B_d^0 is nearly 3 orders of magnitude higher than that for B_c . Again, while the branching fractions for the backgrounds of eqs. (3a) and (3b) are 1.1×10^{-3} and 4.7×10^{-4} respectively [19], that for the signal chain is approximately³ $0.08 \mathcal{B}$ where

$$\mathcal{B} = Br(B_c \rightarrow B_s \ell \nu) + Br(B_c \rightarrow B_s^* \ell \nu) . \quad (4)$$

Using the theoretical prediction of $\mathcal{B} \sim 0.2$ [1], it is easy to see that the background can, *a priori*, be much larger than the signal. However, as Fig. 2 shows, one could still separate the signal from the background by imposing a cut⁴ on the invariant mass for the dilepton pair:

$$m_{\ell\ell} > m_{min} .$$

³Note that this includes the factor of half due to the fact that we consider only the like-sign dilepton case.

⁴The invariant mass for the signal falls far short of the J/ψ and hence such backgrounds can easily be eliminated without losing any of the signal.




Figure 1: *Distribution in the invariant mass of the lepton pair. The solid line corresponds to the signal, while the short- and long-dashed lines refer to backgrounds from the decay chains (3a) and (3b) respectively.*

Clearly, with a strong cut on $m_{\ell\ell}$ most of the background can be eliminated while losing a relatively smaller fraction of the signal. For example, with a m_{min} of 0.8, 1.2 and 1.6 GeV, one retains 64.5%, 33.8% and 11.4% of the signal respectively. For the same set of cuts, the background (3a) gets reduced to 16.7%, 0.86% and 0.001% respectively, while background (3b) falls to 3.4%, 0.01% and 0% of its unrestricted value.

Although a relatively large value of m_{min} would eliminate all of the background, one needs to ensure that the events retained are sufficiently numerous for them to be considered a signal. To make a judicious choice, one thus has to consider the event rates expected. The latter is obviously given by the product $(0.08 \mathcal{B} N \epsilon)$, where N is the number of B_c s produced, and ϵ is the experimental efficiency of detecting the final state. Now, ϵ is the same for both the signal and the background, while the number of B_d s produced can be approximated⁵ to be $500N$.

At the BTeV experiment, $N \simeq 10^8$ even with a conservative estimate. The rate may be slightly higher at the LHC-B. The identification efficiency for each lepton is about 90 %. The D_s is mostly to be recognised through the $\phi\pi$ and K^*K channels, in which the combined branching ratio is 6%. Each of these will give rise to three tracks. Including the trigger efficiency and all kinematic cuts necessary for proper identification, the average efficiency for these three tracks being faithfully reconstructed into a D_s is approximately 2.5%. Thus, with the requirement of proper identification of the other lepton (coming from B_c -decay), the net efficiency of detection of the final states of our concern turns out to be $\epsilon \sim 0.001$. It should be noted that the parent B_c is sufficiently boosted, so that the lepton coming from its decay at the first step stays more or less in the same rapidity interval as the D_s decay tracks and the lepton emanating from B_s -decay.

In Table 2, we present, for a given $N\epsilon$, the number of signal (S) and background (B) events as a function of m_{min} . The significance $S/\sqrt{B}$ (which is proportional to $\sqrt{N\epsilon}$) is then easy to

⁵Approximately only 0.1% of the b quarks produced end up in B_c s while most of them hadronize into B_d and $B^\pm$ with nearly equal probability.




Figure 2: The fraction of the signal and background retained on imposition of a lower bound on the invariant mass of the lepton pair. The solid line corresponds to the signal, while the short- and long-dashed lines refer to backgrounds from the decay chains (3a) and (3b) respectively.

calculate. Since the branching ratios for all other decays in both the signal and the background

m_{min} (GeV)	Signal	Background	
		(3a)	(3b)
0.0	8000 $\mathcal{B}$	55000	23500
0.2	7820 $\mathcal{B}$	49980	19330
0.4	7240 $\mathcal{B}$	36610	10410
0.6	6320 $\mathcal{B}$	21100	3640
0.8	5160 $\mathcal{B}$	9180	813
1.0	3900 $\mathcal{B}$	2780	98
1.2	2700 $\mathcal{B}$	474	3.7
1.4	1770 $\mathcal{B}$	28	0
1.6	910 $\mathcal{B}$	0	0

Table 2: The number of signal and background events retained as a function of m_{min} for $N\epsilon = 10^5$. We assume that 500 as many B_d s are produced as B_c s.

cascades are rather well-known, it should be possible from the above estimates to measure $\mathcal{B}$, the sum of the branching ratios for $B_c \rightarrow B_s^{(*)} l \bar{\nu}$ and match it with the different theoretical predictions. It is interesting to note that, even without imposing any kinematical cuts, it would be possible to have a 5σ signal as long as $\mathcal{B} > 0.17$.

Further distinction of the signal from the backgrounds can be achieved by considering different angular distributions. These distributions in the laboratory frame require the B_c to be boosted appropriately, and one can perform a detailed simulation only by knowing the machine parameters. On the other hand, some interesting features can be observed for the distributions




Figure 3: *The distribution in the angle between the D_s and the slower of the two leptons for different values of the lower bound on the dilepton invariant mass. The solid line corresponds to the signal, while the dashed line refers to the background (the curves for the two backgrounds (3a) and (3b) are virtually indistinguishable).*

Let us, for example, consider, for the signal process $B_c \rightarrow B_s(B_s^*)$, the distribution in the opening angle between the reconstructed D_s and the slower of the two leptons (which, in most of the cases, is the one coming from the primary decay). As shown in figure 3, the distribution is a slowly falling one, as opposed to the corresponding case for the backgrounds, where a peaking for large angle is evident. This can be explained by noting that the background leptons originate from $B_d^0 \rightarrow D_s(D_s^*)D^-$ followed by cascade decay of the D^- . Since the momentum of the lepton in the rest frame of the D^- is smaller than the momentum of the D^- itself, the leptons have trajectories close to that of the D^- . The latter, on the other hand, moves back-to-back with respect to the D_s in the B_c rest frame, making it imperative for the leptons, too, to have a large opening angle against the D_s .

With increasing m_{min} , events with more energetic leptons, or with large opening angles between them, are filtered out. For the signal events, this results in a slightly enhanced likelihood of small opening angle between the slow lepton and the D_s . For the background, on the other hand, an increase in m_{min} does not change this particular distribution to a perceptible extent. This can be traced to the event topology discussed in the last paragraph.

Though the distinctions suggested above are most noticeable in the rest frame of the B_c , they can be utilised if the latter can be reconstructed even at a statistical level. If such reconstruction is possible, the study of the angular distributions will enable one to filter out the signals even with a somewhat relaxed invariant mass cut and thus to be left with a larger number of events.

In conclusion, the study of D_s plus like-sign dilepton signals can serve as extremely useful pointers to the decay $c \rightarrow s$ taking precedence over $b \rightarrow c$ in a B_c meson. Using invariant mass cuts on the dileptons, one can reduce the main backgrounds to a large extent while still preserving a sufficient number of signal events. A careful analysis of such events at the hadronic B-factories can therefore reveal useful information enabling one to ascertain the correctness of different theoretical claims regarding the decay of the charm quark with a b-spectator.

We acknowledge helpful discussions with Tariq Aziz and Sheldon Stone.

References

- [1] See, *e.g.*, S. Gershtein *et al*, hep-ph/9803433.
- [2] F. Abe *et al* (CDF collaboration), *Phys. Rev. Lett.* **81** (1998) 2432; hep-ex/9804014.
- [3] R. Barate *et al* (ALEPH collaboration), *Phys. Lett.* **B402** (1997) 213.
- [4] K. Ackerstaff *et al* (OPAL collaboration), *Phys. Lett.* **B420** (1998) 157.
- [5] P. Abreu *et al* (DELPHI collaboration), *Phys. Lett.* **B398** (1997) 207.
- [6] M. Galdon *et al*, *Nucl. Phys. Proc. Suppl.* **50** (1996) 311.
- [7] E. Eichten and C. Quigg, *Phys. Rev.* **D49** (1994) 5845.
- [8] D. Du, H. Jin and Y. Yang, *Phys. Lett.* **B414** (1997) 130.
- [9] D. Du and Z. Wang, *Phys. Rev.* **D39** (1989) 1342.
- [10] A.Yu. Anisimov *et al*, hep-ph/9809249.
- [11] M. Choi and J.K. Kim, *Phys. Rev.* **D53** (1996) 6670.
- [12] M. Beneke and G. Buchalla, *Phys. Rev.* **D53** (1996) 4991.
- [13] C.H. Chang and Y. Chen, *Phys. Rev.* **D49** (1994) 3399.
- [14] M. Lusignoli and M. Masetti, *Z. Phys* **C51** (1991) 549.
- [15] For a review, see, *e.g.*, S. Stone, hep-ph/9709500.
- [16] E. Braaten, K. Cheung and T. Yuan, *Phys. Rev.* **D48** (1993) R5049.
- [17] A.K. Likhoded *et al* (CDF collaboration), *Nucl. Instrum. Meth.* **A333** (1993) 209.
- [18] M. Wirbel, B. Stech and M. Bauer, *Z. Phys.* **C29** (1985) 637.
- [19] Particle Data Group, C. Caso *et al*, *Euro. Phys. J.* **C3** (1998) 1.